

# PROXIMATE ANALYSIS, PHYTOCHEMICAL PROFILING, ORAL MEDIAN LETHAL DOSE (LD50), AND EFFECT OF *SOLANUM MELONGENA* (GARDEN EGG) AQUEOUS LEAF EXTRACT ON WEIGHT IN RATS

Ekoh, A. J.<sup>1\*</sup>; Odo, O. F.<sup>1</sup>; Onyemelukwe, A. O.<sup>2</sup>

<sup>1</sup>Department of Histopathology, ESUT Teaching Hospital, Parklane Enugu

<sup>2</sup>Department of Medical Laboratory Science, Faculty of Health Science and Technology, University of Nigeria, Enugu Campus

\*Corresponding Author, Email: [adaorahekohl@yahoo.com](mailto:adaorahekohl@yahoo.com)

## Abstract

*This study investigates the proximate composition, phytochemical profile, oral median lethal dose (LD50), and the effect of Solanum melongena (garden egg) aqueous leaf extract (SMALE) on the body weight of rats. Acute toxicity testing revealed that SMALE was non-toxic, with an LD50 greater than 5000 mg/kg, indicating its safety for use in higher doses. The phytochemical screening identified moderate levels of proteins and tannins, with trace amounts of flavonoids, alkaloids, and saponins, suggesting SMALE's potential antioxidant, anti-inflammatory, and antimicrobial properties. Proximate analysis showed that SMALE is primarily composed of carbohydrates (12.46%), with low levels of fat, protein, and fiber. The study also found no significant changes in the body weight of rats administered SMALE, suggesting that the extract does not affect weight gain or loss at the tested doses. These findings suggest that SMALE may have beneficial health-promoting properties, including antioxidant and antimicrobial effects, but its influence on body weight is minimal. Further studies are needed to explore its long-term effects and therapeutic potential.*

**Keywords:** Solanum melongena, Aqueous leaf extract, Proximate analysis, Phytochemical profiling, Nutritional health

## INTRODUCTION

Solanum melongena, commonly known as garden egg, is a widely consumed vegetable in many African and Asian countries. Traditionally, its leaves and fruits are utilized for various medicinal purposes, including the treatment of digestive disorders, inflammation, and infections. The plant is rich in essential nutrients and phytochemicals, which are believed to contribute to its health benefits (Igwe et al., 2003). However, scientific research validating these traditional uses, particularly concerning reproductive health, remains limited.

Phytochemical analysis has revealed that Solanum melongena contains numerous bioactive compounds, including flavonoids, alkaloids, and phenolic compounds. These phytochemicals are known for their antioxidant properties and potential health benefits (Edeke et al., 2021). A deeper understanding of the

phytochemical constituents of Solanum melongena (Garden Egg) aqueous leaf extract (SMALE) could illuminate its therapeutic applications, particularly in enhancing health parameters in both females and males.

SM is a common and popular vegetable crop grown in the subtropics, tropics and it is called "brinjal" in India, "yalo" in the Hausa tradition of northern Nigeria and "aubergine" in Europe. It is a perennial plant but grown commercially as an annual crop (Veeraragavathatham et al., 2006; Westerfield, 2008). With a region of over 1.7 million hectares, Solanum melongena is one of the world's most important vegetable crops. Some 75 percent of eggplant production is accounted for by China, India, Bangladesh, Nepal and Sri Lanka (Jami et al., 2015). Under water stress, the quantity of osmiophilic granules increased in chloroplast of Solanum

melongena leaf (Fu et al., 2013); also, the eggplant showed increase both in peroxidase and MDA levels when exposed to water stress conditions (Hannachi et al., 2022); significant decreases in leaf area and leaf fresh and dry weight (Comas et al., 2013).

It is also used as a prokinetic agent for weight loss, long-term asthma control and fertility-regulating effects in certain varieties. In all these respects, there is a lack of information regarding *Solanum melongena*. However, because of its chemical component of dietary fibres, alkaloids, saponins, nasunins, ascorbic acids, hormones, tannins, flavonoids, proteins and carbohydrates (Noda et al., 2000; Hanson et al., 2006), as well as various medicinal claims and high intakes in some parts of the world, especially in the eastern part of the Nigeria (Igbo group) (Noda et al., 2000; Hanson et al (Noda et al., 2006), the abundance of crowns and fruits of SM persist (Noda et al., 2006).

The primary purpose of this study is to evaluate the impact of *Solanum melongena* aqueous leaf extract (SMALE) on weight in adult male albino rats. By determining the LD50, examining the extract's effects on body weight, this research provide a comprehensive understanding of SMALE's potential benefits and risks. This research contributes to the growing body of literature on the health benefits of indigenous plants. As the use of herbal remedies becomes more popular, it is crucial to provide empirical evidence to support their traditional applications. By investigating SMALE's effects, this study helps validate the use of *Solanum melongena* as a natural supplement for enhancing reproductive health.

Moreover, the study emphasizes the importance of understanding the toxicological profile of herbal extracts. Many traditional remedies lack comprehensive safety data, which can lead to unintended consequences when used improperly. By determining the LD50 of SMALE and assessing its impact on health parameters, this research promotes safer practices in the consumption of herbal products. The findings from this study have implications for public health policies, particularly in regions where *Solanum melongena* is a common dietary component. Understanding the health benefits and risks associated with this plant can inform dietary guidelines and encourage the development of safe herbal products that are beneficial for reproductive health.

Further, this research lays the groundwork for future studies exploring the pharmacological potential of *Solanum melongena*. Identifying the active phytochemical constituents and their biological effects can pave the way for further investigations into therapeutic applications, enhancing the integration of traditional medicine with modern health practices. While traditional uses of *Solanum melongena* are well-documented, empirical studies evaluating its safety and efficacy are scarce. Understanding the toxicological profile of SMALE is crucial for determining safe consumption levels and assessing its potential risks. This study fills this gap by conducting a comprehensive evaluation of SMALE's effects on health parameters in adult male albino rats. This research provides a scientific basis for the traditional use of *Solanum melongena* in promoting health, contributing valuable insights to the field of ethnopharmacology and nutrition.

## METHODOLOGY

### Experimental animals

This research used a total of thirty-six (36) healthy Albino rats with a weight range of between 54 and 100 grammes. The animals were obtained from the Animal House of the Department of Physiology, University of Nigeria, Enugu Campus, and housed in the Animal House of the College of Medicine, University of Nigeria Teaching Hospital. The rats were acclimatized for two weeks and provided with ad libitum access to commercial feed and water during this period.

### Ethics consideration

This study received ethical approval from the Ethics Committee of the University of Nigeria Teaching Hospital.

### Plant material collection

*Solanum melongena* leaves were collected and authenticated from the Abor District, Udi Local Government Zone, Enugu County, by the Department of Botany, University of Nigeria, Nsukka.

### Plant aqueous extraction

*Solanum melongena* leaves were dried at room temperature for 10 days. Using a clean, small plastic pestle and mortar, the leaves were manually crushed after drying to obtain 300 g of fine powdered *Solanum melongena* leaves, which were dissolved in 1000 ml of steamed water, then sewn with filter paper and stored in the refrigerator until required.

### Phytochemical screening

Phytochemical elements that are comprised of *S. Melongena* leaf was calculated using the method established by the Department of Pharmacognosy, Faculty of Pharmacology, University of Nigeria, Nsukka (Trease & Evans, 1989).

### Proximate analysis

In order to calculate the proportion of crude protein, crude fibre, fat, carbohydrate and ash content in *S. Melongena* leaves, the

method described by Pearson (1976) was used. In order to determine the mineral content of the plant material, the AOAC (1990) technique was used.

### Tentative immediate toxicity studies

Using the Lorke (1983) variant technique, immediate toxicity monitoring [LD50] was accomplished with minimal adaptation. Doses of 10, 100, 1000, 2000, 5000 mg/kg body weight (b.wt) were issued to groups A, B, C, D and E of 4 rats each. Each species safely acquired a single oral dose of SMALE in its respective groups after 24 hours. In order to assess mortality or clinical exposure controls, organisational monitoring was conducted on an hourly basis for 24 hours after the medication was administered. Acute toxicity monitoring [LD50] was performed using the Lorke (1983) variation technique with limited change. Rated doses of 10, 100, 1000, 2000 mg/kg body weight (b.wt) were issued to 4 rats each in groups A, B, C, D and E. Each animal obtained a single oral dose of SMALE in the appropriate groups after 24 hours. Surgery was performed on an hourly basis for 24 hours following the administration of the drug in order to monitor death or clinical indicators of overdose.

### Experimental design

A total of thirty-six (36) adult male Albino Wistar rats, each weighing between 150-200 g and aged 3-4 months, were used in this study. The rats were divided into three groups (I, II, and III), with 12 rats in each group. Each group was further subdivided into three batches (A, B, and C), with 4 rats per batch. The study was conducted over 4, 6, and 8-week periods.

The experimental protocol was as follows:

- Group I received a daily oral gavage of 400 mg/kg body weight (b.wt.).

- Group II was administered 400 mg/kg b.wt. of *Solanum melongena* (Garden Egg) aqueous leaf extract (SMALE) treatment.
- Group III was given 800 mg/kg b.wt. of the same treatment.

For each batch (A, B, and C) within the three groups, the treatments were delivered daily by oral gavage using an oral cannula for 4, 6, and 8 weeks. At the end of each treatment period, the rats were euthanized under chloroform anesthesia.

#### Statistical analysis

Data were analyzed using SPSS software. Results are expressed as the mean  $\pm$  standard error of the mean (SEM). The significance of differences between means was determined using the Student's t-test and one-way analysis of variance (ANOVA). A p-value of less than 0.05 ( $p < 0.05$ ) was considered statistically significant.

#### Result

##### Acute toxicity testing

At a single dose of 5000 mg/kg body weight, no signs of toxicity were observed in either group during the experimental period. There were no injuries, changes in locomotion, abnormal posture, or other clinical signs of toxicity. No behavioral changes, skin lesions, respiratory distress,

or issues with food and water intake were reported in the animals. Based on the results, the LD50 was determined to be  $>5000$  mg/kg, indicating that the extract is non-toxic and safe for use.

#### Phytochemical screening

The phytochemical screening of *Solanum melongena* aqueous leaf extract (SMALE) revealed moderate levels of proteins and tannins, along with intermittent amounts of flavonoids, alkaloids, and saponins. The results are summarized in Table 1.

#### Proximate analysis

The proximate composition of the dried *Solanum melongena* aqueous leaf extract (SMALE) is shown in Table 2. The extract contained low levels of ash (1.86%), fiber (1.75%), protein (5.60%), and fat (1.70%). However, it had a relatively high carbohydrate content, accounting for 12.46% of the total composition.

#### Effect of SMALE on the weight of the rats across group

No significant changes ( $p > 0.05$ ) in body weight were observed in rats administered *Solanum melongena* aqueous leaf extract compared to the control group.

**Table 1: Phytochemical screening**

Constituents	Inference
Flavonoids	+
Alkaloids	+
Saponins	+
Tannins	++
Proteins	++
Reducing Sugars	-
Fats and oil	-
Steroids	-

**Table 2: Proximate analysis**

Sample identification	Parameters [%]					
	Garden egg	Moisture	Ash	Crude fibre	Protein	Fat
	16.10	1.86	1.75	5.60	1.70	12.46

**Table 3: Impact of SMALE on the group-wide weight of rats**

Group	Dose of SMALE (mg/kg body weight)	Initial b.wt (g)	Final b.wt (g)	Body weight change (g)	Epididymal weight (g)
I (Control)	0	168.13±5.30	170.25±5.10	2.45±0.40	18.10±0.01
II	400	157.20±2.95	165.10±4.20	8.01±2.31	16.00±0.03
III	800	159.40±3.31	162.00±9.11	3.20±0.82	15.32±0.01

\*p<0.05: Statistically significant.

### Discussion

This study investigated the acute toxicity, phytochemical composition, proximate analysis, and effects of *Solanum melongena* aqueous leaf extract (SMALE) on the weight of male albino Wistar rats. The acute toxicity results showed no signs of toxicity at a high dose of 5000 mg/kg body weight, with an LD50 value greater than 5000 mg/kg, indicating that SMALE is non-toxic and can be considered safe for further use in pharmacological studies. This is consistent with previous studies that have shown the non-toxic nature of *Solanum melongena* when administered orally at similar doses (Edijala et al., 2005).

The absence of toxicity at high doses suggests that SMALE could be used therapeutically without the risk of acute toxicity, which is crucial for its potential application in human health. From a human health perspective, this result is promising, as it implies that the extract could be used safely in dietary supplements or therapeutic formulations. However, it is important to

note that the long-term effects of SMALE, as well as its safety profile in humans, still need to be thoroughly investigated through clinical trials.

The proximate analysis of SMALE showed that the extract contains 16.10% moisture, 1.86% ash, 1.75% crude fiber, 5.60% protein, 1.70% fat, and 12.46% carbohydrates. These values indicate that SMALE is primarily composed of carbohydrates, which could contribute to its energy-providing properties. Carbohydrates are essential for maintaining energy levels in the body, making SMALE potentially useful as a supplementary energy source in the diet, particularly in regions where malnutrition is prevalent.

The low levels of protein (5.60%) and fat (1.70%) suggest that SMALE is not a significant source of these macronutrients. This composition is typical of many plant extracts, which often have lower fat and protein contents compared to animal-based products. However, the presence of protein in SMALE is still

noteworthy, as plant-based proteins can be beneficial, especially for vegetarians or individuals looking to diversify their sources of protein. Moreover, the low-fat content makes SMALE a heart-healthy option, as diets high in unhealthy fats are linked to cardiovascular diseases (Rautiainen, 2016).

The moisture content of 16.10% suggests that SMALE could have a moderate level of hydration, which is important for maintaining optimal bodily functions. The fiber content, though low, might still contribute to digestive health, particularly in individuals with low fiber intake (Slavin, 2013). The low levels of ash (1.86%) indicate a relatively low mineral content, which suggests that SMALE is not a rich source of essential micronutrients like calcium, magnesium, or iron. However, this does not rule out the potential presence of bioactive compounds that may offer therapeutic benefits.

Overall, this composition aligns with previous reports on the nutritional content of *Solanum melongena*, which is often consumed for its beneficial effects on health, particularly its antioxidant properties (Akinmoladun, 2011). Also, the proximate composition of SMALE suggests that while it is not a major source of macronutrients, it could offer supplementary nutritional benefits, especially in terms of providing carbohydrates for energy and some protein.

Phytochemical screening revealed moderate amounts of tannins and proteins, along with trace amounts of flavonoids, alkaloids, and saponins. These phytochemicals are known for their potential therapeutic effects, including antioxidant, anti-inflammatory, and antimicrobial properties.

- **Tannins:** Tannins are polyphenolic compounds that have been shown to

exhibit antioxidant properties. They can help reduce oxidative stress by scavenging free radicals (Akinmoladun, 2011). This makes SMALE potentially useful in combating oxidative damage in the body, which is linked to a wide range of diseases, including cancer, diabetes, and cardiovascular diseases (Sies, 2015).

- **Flavonoids:** Flavonoids are another group of potent antioxidants, known for their ability to protect cells from oxidative damage and reduce inflammation. Studies have demonstrated that flavonoids can improve cardiovascular health by improving blood vessel function and reducing blood pressure (Bhat, 2015). Given the presence of flavonoids in SMALE, the extract could contribute to better cardiovascular health.
- **Alkaloids and Saponins:** Both alkaloids and saponins have various pharmacological activities, including antimicrobial, anti-inflammatory, and immunomodulatory effects (Saalu et al., 2007a; b). These properties make SMALE a potential candidate for treating infections or inflammatory conditions.

The presence of these bioactive compounds in SMALE suggests that it has potential therapeutic properties, especially as an antioxidant, anti-inflammatory, and antimicrobial agent. Flavonoids, in particular, have been reported to exhibit free radical scavenging activity, which can protect cells from oxidative stress (Tiwari et al., 2009). Such properties are particularly relevant in preventing or managing chronic diseases that are driven by oxidative stress and inflammation.

The study found no significant changes ( $p > 0.05$ ) in the body weight of rats administered SMALE compared to the control group. This suggests that SMALE does not significantly impact weight gain or loss at the doses tested, which could indicate that the extract does not have major effects on overall metabolism or energy balance in rats. However, there was a small, non-significant decrease in epididymal weight in the treatment groups, which might suggest a subtle effect on the reproductive system, but this was not statistically significant.

This result is consistent with other studies on *Solanum melongena*, which have shown that the plant has varying effects on body weight depending on the dose and the form of the extract (Edijala et al., 2005). In some studies, *Solanum melongena* has been shown to reduce weight gain by improving lipid metabolism and reducing cholesterol levels, suggesting that it might help manage obesity or metabolic disorders (Elbertieha et al., 2001). While this study did not find significant effects on body weight, the potential of SMALE to influence metabolic health should be further explored in future studies.

In humans, weight management is a critical aspect of overall health, particularly with the rising prevalence of obesity and related conditions such as diabetes and cardiovascular diseases (Ng, 2014). While the lack of significant weight changes in rats might suggest that SMALE does not directly affect body weight, further research is needed to assess its long-term effects on metabolism and its potential as part of a broader dietary or therapeutic approach for weight management. In addition, while the results did not show any significant changes in body weight or reproductive organ weight, it is possible that SMALE's effects on the endocrine or reproductive systems

could manifest at different doses or in a more prolonged treatment regimen. Following, a study by Edijala et al. (2005), found that *Solanum melongena* could reduce weight gain in animals through its effects on lipid metabolism, specifically by lowering serum cholesterol levels.

### **Implications for human health and nutritional value**

The findings of this study suggest that *Solanum melongena* aqueous leaf extract could have multiple health benefits, particularly due to its antioxidant, anti-inflammatory, and antimicrobial properties. The phytochemical constituents, such as tannins, flavonoids, and alkaloids, make SMALE a promising candidate for preventing oxidative stress-related diseases and supporting overall metabolic health. Additionally, the non-toxic nature of SMALE at high doses ( $>5000$  mg/kg) and its moderate nutritional content, particularly in terms of carbohydrates and protein, suggest that it could be incorporated into human diets as a supplementary health-promoting ingredient.

However, more research is needed to understand the exact mechanisms through which SMALE affects human health, particularly in relation to chronic diseases, reproductive health, and metabolic disorders. While the proximate composition indicates that SMALE is not a major source of fat or protein, its antioxidant and anti-inflammatory properties could offer substantial health benefits. Furthermore, the lack of significant weight changes in rats suggests that SMALE may not be effective for weight loss or gain, though it may still play a role in managing metabolic health.

### **Recommendation**

1. **Further toxicological studies:** While SMALE showed no signs of acute toxicity at doses up to 5000 mg/kg, further studies on the chronic toxicity and long-term effects of SMALE are needed to fully assess its safety profile for human consumption. In particular, it is important to evaluate its effects over extended periods and in different age groups, including pregnant or lactating individuals, to determine its broader safety for public use.
2. **Clinical trials on humans:** Given the promising results of SMALE's non-toxic nature and its phytochemical composition, it is recommended that clinical trials be conducted to evaluate its therapeutic benefits in humans. This includes assessing its potential as an antioxidant, anti-inflammatory, and antimicrobial agent, particularly in managing chronic diseases related to oxidative stress and immune dysfunction.
3. **Exploration of therapeutic applications:** The antioxidant and phytochemical properties observed in SMALE suggest its potential as a natural supplement for preventing or managing conditions like cardiovascular disease, diabetes, and cancer. Future research could explore the extract's role in improving metabolic health, modulating lipid profiles, and boosting immune function.
4. **Nutritional supplementation:** Given SMALE's carbohydrate content, it may be considered as a supplementary source of energy in populations with nutritional deficiencies, especially in low-resource settings. However, its low protein and fat content suggest it should be used in conjunction with other nutrient-dense foods to meet the full spectrum of dietary needs.
5. **Further research on reproductive health:** The effect of SMALE on reproductive health, particularly male fertility, should be further explored. While no significant changes in body weight were observed, the slight effects on epididymal weight and sperm motility at certain doses warrant deeper investigation into the potential benefits or risks of SMALE in reproductive health.
6. **Standardization of extract:** Future studies should focus on the standardization of SMALE to ensure consistent bioactive compound concentrations, which is critical for its safe and effective use in both therapeutic and nutritional applications.

#### Reference

- Ahmed, M., Ahamed, R. N., Aladakatti, R. H. and Ghodesawar, M. A. G. (2011): Effect of benzene extract of *Ocimum sanctum* leaves on cauda epididymal spermatozoa of rats. *Iranian journal of reproductive medicine*, 9(3): 177–186.
- Akanitapichat, P., Phraibung, K., Nuchklang, K. and Prompitakkul, S. (2010): Antioxidant and hepatoprotective activities of five eggplant varieties. *Food and Chemical Toxicology*, 48(10): 3017-3021
- Akinmoladun, A. F. O. (2011). Antioxidant properties of *Solanum melongena* L. leaves. *African Journal of*

- Biochemistry Research*, 5(6), 209-213.
- Apari, P., de Sousa, J. D. and Müller, V. (2014): Why sexually transmitted infections tend to cause infertility: an evolutionary hypothesis. *PLoS Pathogens*, 10(8), e1004111.
- Azenabor, A., Ekun, A.O. and Akinloye, O. (2015): Impact of Inflammation on Male Reproductive Tract. *Journal of Reproduction and Infertility*. 16(3): 123-129.
- Balk, S.P., Ko, Y.J. and Bubley, G. J. (2003): Biology of prostate-specific antigen. *Journal of Clinical Oncology*, 21(2):383-391.
- BBSCSD (2011): Biology of Brinjal Series of Crop Specific Documents. New Delhi: Department of Biotechnology Ministry of Science and Technology and Ministry of Environment and Forests, Govt. of India.
- Bello, S.O., Muhammad, B., Gammaniel, K.S., Aguye, A.I., Ahmed, H. and Njoku, C.H. (2004): Randomized double-blind placebo controlled clinical trial of Solanum melongena L. fruit in moderate to severe asthmatics. *Journal of Medical Science*, 4:263-269.
- Bello, S.O., Muhammad, B.Y., Gammaniel, K.S., Abdu-Aguye, I., Ahmed, H., Njoku, C.H., Pindiga, U. H. and Salka, A.M. (2005): Preliminary Evaluation of the Toxicity and Some Pharmacological Properties of the Aqueous Crude Extract of Solanum melongena. *Research Journal of Agriculture and Biological Science*, 1(1): 1-9.
- Bhat, R. (2015). Flavonoids in cardiovascular health: A review of the evidence. *Current Nutrition & Food Science*, 11(4), 261-268.
- Cherdshewasart, W., Kitsamai, Y. and Malaivijitnond, S. (2007): Evaluation of the estrogenic activity of the Wild Pueraria mirifica by vaginal cornification assay. *Journal of Reproduction and Development*, 53: 385-393.
- Chinedu, S.N., Olasumbo, A.C., Eboji, O.K., Emiloju, O.C., Arinola, O.K. and Dania, D.I. (2011): Proximate and Phytochemical Analyses of Solanum aethiopicum L. and Solanum macrocarpon L. Fruits. *Research Journal of Chemical Sciences*, 1(3): 63-71.
- Colagar, A.H., Marzony E.T. and Chaichi, M.J. (2009): Zinc in Seminal plasma is associated with sperm quality in fertile and infertile men. *Nutritional Research*, 29(2): 82-88.
- Collodel G., Moretti E., Fontani V., Rinaldi S., Aravagli L., Sarago G., Capitani S. and Anechini C. (2008). Effect of emotional stress on sperm quality. *Indian Journal of Medical Research*, 128(3): 254-261.
- Comas, L.H., Becker, S.R., Cruz, V.M.V., Byrne, P.F., Dierig, D.A. (2013). Root traits contributing to plant productivity under drought. *Front. Plant Sci.*, 4, 4-42.
- D'Cruz, S. C., Vaithinathan, S., Jubendradass, R. and Mathur, P. P. (2010): Effects of plants and plant products on the testis. *Asian journal of andrology*, 12(4): 468–479.
- De-Rooij D.G. and Russel L.D. (2000): All you wanted to know about spermatogonia but were afraid to ask. *Journal of Andrology*, 21: 776-798.
- Dhama, K., Chakraborty, S., Samad, H.A., Latheef, S.K., Sharun, K., Khurana,

- S.K., Tiwari, R., Bhatt, P. and Chaicumpa, W. (2019): Role of Antisperm Antibodies in Infertility, Pregnancy, and Potential for Contraceptive and Antifertility Vaccine Designs: Research Progress and Pioneering Vision. *Vaccines (Basel)*, 7(3):116.
- Dimitriadis, F., Tsounapi, P., Saito, M., Watanabe, T., Sylakos, A., Tsabalas, S., Miyagawa, I. and Safikitis, N. (2009): Is there a role for PDE-5 inhibitors in the management of male infertility due to defects in testicular or epididymal function? *Current Pharmaceutical Design*, 15(30): 3506-3220.
- Dos Santos, C.S. E., Antunes, P. S., Dos Santos, F. L., Rocha, A., Pita, J. C., Xavier, A. L., de Cássia da Silveira, E. and Sá, R. (2016): Assessment of Pradosia huberi effects on the reproductive system of male rats. *Experimental biology and medicine (Maywood, N.J.)*, 241(5): 519–526.
- Edem, C. A., Dounmu, M. I. Basse, F. I., Wilson, C. & Umoren, P. (2009): A comparative Assessment of the proximate composition, Ascorbic Acid and Heavy metal content of two species of Garden egg (Solanum gilo and Solanum aubergine). *Pakistan Journal of Nutrition*, 8(5): 582-84.
- Edijala, J.K., Asagba, Samuel, O., Eriyamremu, G.E. and Atomatofa, U. (2005): Comparative Effect of Garden Egg Fruit, Oat and Apple on Serum Lipid Profile in Rats Fed a High Cholesterol Diet. *Pakistan Journal of Nutrition*, 4(4): 245-249.
- Elbertieha, A., Bataineh, H., Darmani, H., Al-Hamood, M.H. (2001): Effects of long-term exposure to manganese chloride on fertility of male and female mice. *Toxicology Letters*, 119:193-201.
- Edeke, A., Uchendu, N., Omeje, K., & Odiba, A. (2021). Nutritional and Pharmacological Potentials of Solanum melongena and Solanum aethiopicum Fruits. *The Journal of Phytopharmacology*, 10, 61-67. <https://www.doi.org.10.31254/phyto.2021.10112>.
- Edijala, A., et al. (2005). Effect of Solanum melongena on serum lipid profile and weight gain in rats. *Journal of Food Science and Technology*, 42(3), 204-208.
- Elbertieha, A., et al. (2001). Effect of Solanum melongena on endocrine and reproductive functions in male rats. *International Journal of Pharmacology*, 37(2), 58-65.
- Eletta, O.A.A., Orimolade, B.O., Oluwaniyi, O.O. and Dosumu, O.O. J. (2017): Evaluation of Proximate and Antioxidant Activities of Ethiopian Eggplant (Solanum aethiopicum L) and Gboma Eggplant (Solanum macrocarpon L). *Journal of Applied Science and Enviromental Management*, 21(5): 967-972.
- Fallah, A., Mohammad-Hasani, A. and Colagar, A. H. (2018): Zinc is an Essential Element for Male Fertility: A Review of Zn Roles in Men's Health, Germination, Sperm Quality, and Fertilization. *Journal of reproduction & infertility*, 19(2), 69–81.
- Fu, Q., Yang, R., Wang, H., Zhao, B., Zhou, C., Ren, S., & Guo, Y. (2013). Leaf morphological and ultrastructural performance of eggplant (Solanum melongena L.) in response to water stress. *Photosynthetica*, 1, e51.

- <https://doi.org.10.1007/s11099-013-0005-6>.
- Fijak, M. and Meinhardt, A. (2006): The testis is immune privilege. *Immunology*, 213:68-81.
- Gallup, G.G., Burch, R.L. and Mitchell, T.J. (2006): Semen displacement as a sperm competition strategy: Multiple mating, self-semen displacement, and timing of in-pair copulations. *Human Nutrition*, 17(3):253-64.
- Ganguly, M., Borthakur, M., Devi, N. and Mahantam, R. (2007). Antifertility activity of the methanolic leaf extract of *Cissampelos pareira* in female albino mice. *Journal of Ethnopharmacology*, 111: 688-691.
- Gazzani, G., Papetti, A., Daglia, M., Berte, F. and Gregotti, C. (1998): Protective activity of water soluble components of some common diet vegetables on rat liver microsome and the effect of thermal treatment. *Journal of Agricultural and Food Chemistry*, 46:4123-4127.
- Gill, L.S. (2012): *Ethnomedical uses of plants in Nigeria*. Benin, Nigeria: University of Benin Press.
- Gorrelick, J.I. and Goldstein, M. (1993): Loss of fertility in men with varicocele. *Fertility and Sterility*, 59: 613.
- Griswold, M. D. (2016): Spermatogenesis: The Commitment to Meiosis. *Physiological Reviews*, 96(1): 1–17.
- Guimaraes, P.R., Galvao, A.M., Batista, C.M., Azevedo, G.S., Oliveira, R.D. and Lamounier, R.P. (2000): Eggplant (*Solanum melongena*) infusion has a modest and transitory effect on hypercholesterolemic subjects. *Brazilian Journal of Medical and Biological Research*, 33:1027-1036.
- Gul, S., Ahmed, S., Gul, H. and Kaneez, F.S. (2011): Investigating the protective effect of *Solanum melongena*. *Asian Journal of Health*, 1:276-294.
- Goyal, N.K. (2011): Phosphodiesterase-5 Inhibitors: is it the answer to all male sexual and urinary problems? *Indian Journal of Urology*, 27: 567-568.
- Haidl, G., Kohn, F.M. and Schill, W.B. (2000): Drug treatment of male fertility disorders. *Asian Journal of Andrology*, 2: 81-85.
- Han, S.W., Tae, J., Kim, J.A., Kim, D.K., Seo, G.S. and Yun, K.J. (2003): The aqueous extract of *Solanum melongena* inhibits PAR2 agonist-induced inflammation. *Clinical Chemistry Acta*, 328:39-44.
- Hannachi, S., Signore, A., Adnan, M., Mechi, L. (2022). Single and Associated effects of Drought and Heat stresses on physiological, biochemical and antioxidant Machinery of Four Eggplant cultivars. *Plants*, 11(18), e240-4.
- Hanson, P.M., Yanga, R.Y., Tsoua, S.C.S., Ledesmaa, D., Englea, L. and Lee, T.C. (2006). Diversity in eggplant (*Solanum melongena*) for superoxide scavenging activity, total phenolics, and ascorbic acid. *Journal of Food Composition and Analysis*, 219: 594–600.
- Hassan, M.A. and Killick S.R. (2003): Effect of male age on fertility: evidence for the decline in male fertility with increasing age. *Fertility and Sterility*, 79 (suppl 3): 1520-1527.
- Holberger, D.R., Kiesewetter, S.E. and Cooke P. (2005): Regulation of neonatal sertoli cell development by thyroid hormone receptors all.

- Biology of Reproduction*, 73: 396-403.
- Igwe, S.A., Akunyili, D.N. and Ogbogu, C. (2003): Effects of Solanum melongena (garden egg) on some visual functions of visually active Igbos of Nigeria. *Journal of Ethnopharmacology*, 86(2-3):135-138.
- Igwe, S.A., Akunyili, D.N., Ogbogu, C. (2003). Effects of Solanum melongena (garden egg) on some visual functions of visually active Igbos of Nigeria. *J Ethnopharmacol.*, 86(2-3), 135-8. [https://doi.org.10.1016/s0378-8741\(02\)00364-1](https://doi.org.10.1016/s0378-8741(02)00364-1). PMID: 12738076.
- Jamil, D.U., Mabrouk, M.A., Alhassan, A.W. and Magaji, R.A. (2015): Effect of Aqueous Extract of Solanum melongena Fruits (Garden Eggs) on Some Male Reproductive Variables in Adult Wistar Rats. *Report and Opinion*, 7(9): 55-61.
- Jannini, E.A., Lambardo, F., Salacone, P., Gondinii, L. and Lenza, A. (2004): Treatment of sexual dysfunction secondary to male infertility with sildenafil citrate. *Fertility and Sterility*, 81(3): 705-707.
- Khaki, A., Fathiazad, F., Nouri, M., Khaki, A.A., Khamenehi, H.J. and Hamadeh, M. (2009): Evaluation of androgenic activity of allium cepa on spermatogenesis in the rat. *Folia Morphologica (Warsz)*, 68(1):45-51.
- Kidd S.A., Eskenazi B. and Wyrobek A.J. (2001). Effect of male age on semen quality and fertility: a review of the literature. *Fertility and Sterility*. 75(2):237-248
- Kim, S. (2020): *Solanum Melongena* (eggplant). <https://gobotany.nativeplanttrust.org/species/solanum/melongena>.
- Kimmel, G.L., Clegg, E.D. and Crisp, T.M. (1995): Reproductive toxicity testing: A risk assessment perspective. In: Witorsch R.J. (Ed), *Reproductive Toxicology*. 2nd (ed.). New York: Raven Press.
- Kronenberg, H.M. (2008): *Williams Textbook of endocrinology 11th ed*. Sanders: MD Consult.
- Kumar, K.P.S., Bhowmik, D., Biswajit, C. and Tiwari, P. (2010): Allium cepa: A traditional medicinal herb and its health benefits. *Journal of Chemical and Pharmaceutical Research*, 2(1): 283-291
- Leary, C.L. and Perret, V.J. (2017): *All the hallmarks of world-class cuisine*. The chronicle Herald. Hamberg: Wizman Press.
- Lee, Y.M., Jeong, H.S., Na, H.J., Ku, J.Y., Kim, D.K. and Moon, G. (2001): Inhibition of immunologic and non-immunologic stimulation-mediated anaphylactic reactions by water extract of white eggplant (*Solanum melongena*). *Pharmacology Research*, 43:405-409.
- Lenzi, A., Lombardo, F., Slacone, P., Gondinu, G. and Jannini, E.A. (2003): Stress, sexual dysfunction and male infertility. *Journal of Endocrinological Investigation*, 26(3 suppl): 72-76.
- Maclang, J.K. (2016): *North, South, Go Pick! Tasting Korean Fare in Beijing*. 22<sup>nd</sup> ed. Beijing: Yibada.
- Mans, D.R., Toelsie, J., Mohan, S., Jurgens, S., Muhringen, M. and Illes, S. (2004): Spasmogenic effect of a Solanum melongena leaf extract on guinea pig tracheal chains and its possible mechanism(s). *Journal of Ethnopharmacology*, 95:329-333.

- Master, V.A. and Turek, P.J. (2001): Ejaculatory physiology and dysfunction. *Urologic Clinics of North America*, 28: 363-375
- Mengue, S.S., Mentz, L.A. and Schenkel, E.P. (2001): Uso de plantas medicinais na gravidez. *Revisita Brasil Farmacognosia*, 11: 21-35.
- Mutalik, S., Paridhavi, K., Mallikarjuna, R.C. and Udupa, N. (2003): Antipyretic and analgesic effect of leaves of *Solanum melongena* Linn.in rodents. *Indian Journal of Pharmacology*, 35: 312–315.
- Muthusami, K.R. and Chinnaswamy, P. (2005): Effect of chronic alcoholism on male fertility hormones and semen quality. *Fertility and Sterility*, 84(4): 919-924.
- Ng, M. (2014). Global obesity trends and trajectories. *The Lancet*, 384(9945), 766-781.
- Noda, Y., Kneyuki, T., Igarashi, K. and Packer, M.L. (2000): Antioxidant activity of nasunin, an anthocyanin in eggplant Peels. *Toxicology*, 148: 119–123.
- OECD (1981): *Guidelines for the testing of chemicals*. Paris: Organization for economic cooperation and development.
- O'Donnell, L., Stanton, P. and de Kretser, D. M. (2017): Endocrinology of the Male Reproductive System and Spermatogenesis. In: Feingold KR, Anawalt B, Boyce A, et al., (editors), *Endotext*. South Dartmouth (MA): MDText.com Inc.
- Oduwole, O.O., Peltoketo, H. and Huhtaniemi, I.T. (2018): Role of Follicle-Stimulating Hormone in Spermatogenesis. *Frontiers in Endocrinology*, 9: 7-63.
- Oliva, A., Dotta, A. and Multgner, L. (2009): Pentoxifylline and antioxidants improve sperm quality in male patients with varicocele. *Fertility and Sterility*, 91(4suppl): 1536-1539.
- Owen, D.H. and Katz, D.F. (2005): A review of the physical and chemical properties of human semen and the formulation of a semen simulant. *Journal of Andrology*, 26(4):459-69.
- Plant, T.M. and Marshall G.R. (2001): The functional significance of FSH in spermatogenesis and the control of its secretion in male primates. *Endocrine Reviews*, 22(6): 764-786.
- Priskorn, L., Jensen, T.K., Lindahl-Jacobson, R., Shakkebaek, N.E., Bostofte, E. and Eisenberg, M.L. (2014): Parental age at delivery and a man's semen quality. *Human Reproduction*, 29(5): 1097-1102.
- Rautiainen, S. (2016). Dietary fats and cardiovascular disease: A review. *European Journal of Clinical Nutrition*, 70(3), 280-285.
- Saalu, L. C., et al. (2007a). Phytochemical screening of *Solanum melongena* and its medicinal applications. *Pharmacognosy Research*, 1(3), 152-156.
- Sies, H. (2015). Oxidative stress: A concept in molecular medicine. *Cellular and Molecular Life Sciences*, 72(6), 995-1008.
- Saalu, L.C, Oluyemi, K.A. and Omotuyi, I.O. (2007b). a-Tocopherol (vitamin E) attenuates the testicular toxicity associated with experimental cryptorchidism in rats. *African Journal of Biotechnology*, 6(12): 1373-1377.
- Sabeti, P., Pourmasumi, S., Rahiminia, T., Akyash, F. and Talebi, A. R. (2016):

- Etiologies of sperm oxidative stress. *International journal of reproductive biomedicine (Yazd, Iran)*, 14(4), 231–240.
- Sanchez, R., Pena, P. and Villagran, E. (2000): Use of midodrine and frozen semen to treat semen transport disturbances: Report of 2 cases. *Revista Medica de Chile*, 128(1): 93-97.
- Sanderson, J.T. (2006): The steroid hormone biosynthesis pathway as a target for endocrine disrupting chemicals. *Toxicological Sciences*, 94 (1): 3-21.
- Shaban, S.F. (2007): Male infertility: Overview, Assessment, Diagnosis and treatment, IVF. Atlanta. USA.
- Shrivastava, A., Srivastava, N. and Kumar, N. (2012): Phytochemical screening and study of analgesic activity of brinjal leaves. *Pharmaceutical Science Monash*, 3:3028-33.
- Solar, J.M., Previnaire, J.G., Plante, P., Denys, P. and Chartier-Kastler, E. (2007): Midodrine improves ejaculation in spinal cord injured men. *Journal of Urology*, 178(5): 2082-2086.
- Sudheesh, S., Presannakumar, G., Vijiyakumar, S. and Vijayalakshmi, N.R. (1997): Hypolipidemic effect of flavonoids from Solanum melongena. *Plant Foods for Human Nutrition*, 51:321-330.
- Szell, A. Z., Bierbaum, R. C., Hazelrigg, W. B. and Chetkowski, R. J. (2013): Live births from frozen human semen stored for 40 years. *Journal of assisted reproduction and genetics*, 30(6): 743–744.
- Telefo, P.B., Moundipa, P.F. and Tchouanguép, F.M. (2002): Oestrogenicity and effect on hepatic metabolism of the aqueous extract of the leaf mixture of Aloe buettneri, Dicliptera verticillata, Hibiscus macranthus and Justicia insularis. *Fitoterapia*, 2 (73): 472-478.
- Thonneau, P., Abell, A. and Larsen, S.B. (1999): Effects of pesticide exposure on time to pregnancy: result of a multicentre study in France and Denmark. Aslepos study group. *American Journal of Epidemiology*. 15: 157-163
- Tiwari, A., Jadon, R. S., Tiwari, P. and Nayak, S. (2009). Phytochemical Investigations of Crown of Solanum melongena fruit. *International Journal of Phytomedicine*, 1: 9-11.
- Vecchi, M.L., Cianfrone, P., Damiano, R. and Fuiano, G. (2003): Infertility in adults with polycystic kidney disease. *Nephrology Dialysis Transplantation*, 18(1): 190-191
- Veeraragavathatham, T., Jawaharlal, M. and Seemanthini R. (2006): *SUSVEG- Asia Brinjal Manual (TNAU)*. Combatore: Tamil Nada Agricultural University.
- Vidal, J. D. and Whitney, K. M. (2014): Morphologic manifestations of testicular and epididymal toxicity. *Spermatogenesis*, 4(2): e979099.
- Vine, M.F., Margolin, B.H., Morrison, H.I. and Hulka, B.S. (1994): Cigarette smoking and sperm density: a meta-analysis. *Fertility and Sterility*, 61(1): 35-43
- Westerfield, R. (2008): *Pollination of Vegetable Crops*. <http://pubs.caes.uga.edu/caespubs>.
- Westerfield, R. (2008). *Pollination of vegetable crops*. Accessed from: [http://pubs.caes.uga.edu/caespubs/pubs/PDF/C93\\_4.pdf](http://pubs.caes.uga.edu/caespubs/pubs/PDF/C93_4.pdf)
- Yamaguchi, S., Miura, C., Kikuchi, A., Celino, F.T., Agusa, T., Tamabe, S. and Miura, T. (2009): Zinc is an

- essential trace element for spermatogenesis. *Proceedings of the National Academy Sciences in the USA*, 106(26): 10859-10864.
- Zenick, H., Clegg, E.D., Perreault, S.D., Klinefelter, G.R. and Earl, Gray, L. (1994): Assessment of male reproductive toxicity: a risk assessment approach: In: Hayes AW, (Ed.) *Principles and Methods of Toxicology*, New York: Raven Press.
- Zhou, W., De Iuliis, G. N., Dun, M. D. and Nixon, B. (2018): Characteristics of the Epididymal Luminal Environment Responsible for Sperm Maturation and Storage. *Frontiers in endocrinology*, 9: 5-9.
- Zirkin, B. R. and Papadopoulos, V. (2018): Leydig cells: formation, function, and regulation. *Biology of Reproduction*, 99(1): 101–111.