

HISTOLOGICAL EFFECTS ON REPRODUCTIVE ORGANS AND CAUDA EPIDIDYMAL SPERM COUNT AND MOTILITY: EFFECT OF SOLANUM MELONGENA (GARDEN EGG) AQUEOUS LEAF EXTRACT [SMALE] IN ADULT ALBINO RATS

Ekoh, A. J.^{1*}; Onyemelukwe, A. O.²; Odo, O. F.¹

¹Department of Histopathology, ESUT Teaching Hospital, Parklane Enugu

²Medical Laboratory Science, Faculty of Health Science and Technology, University of Nigeria, Enugu Campus

*Corresponding Author, Email: adaorahekoh1@yahoo.com

Abstract

This study investigates the histological effects of Solanum melongena (garden egg) aqueous leaf extract (SMALE) on the reproductive organs, specifically focusing on sperm motility, sperm count, and histological changes in the testes and epididymis of adult male albino rats. The rats were administered varying doses of SMALE (400 mg/kg and 800 mg/kg body weight) over a period of time. Sperm motility and count were assessed, with significant improvements observed in the 400 mg/kg dose group, indicating a dose-dependent enhancement in sperm function. Histological examination revealed normal testicular architecture in the control group, with intact spermatocytes and spermatids. The 400 mg/kg SMALE treatment showed increased proliferation and maturation of spermatocytes and Leydig cells, while the 800 mg/kg dose led to signs of testicular degeneration, characterized by atrophied Leydig cells and pyrosis. These findings suggest that SMALE at moderate doses may enhance reproductive health, while higher doses could induce toxicity and impair sperm maturation. The study highlights the potential of Solanum melongena as a natural supplement for male fertility, with important implications for reproductive health and safety.

Keywords: Solanum melongena, Fertility, Phytochemicals, Antioxidants.

INTRODUCTION

Globally, medicinal plants have long been utilized to treat various diseases; however, some plant compounds may have toxic effects on the body. Certain plants are known to possess teratogenic and abortifacient properties, while others are thought to negatively impact the male reproductive system (Mengue et al., 2001; Dos Santos et al., 2016). The male reproductive system can be adversely affected by a range of factors, including interference with sexual maturation, sperm development and transport, sexual behavior, and overall reproductive function (Kimmel et al., 1999). These effects can result from substances that either disrupt or promote the production of testosterone, a key hormone in male reproductive health (D'Cruz et al., 2010). Some studies have shown that toxic agents can impair the function of the epididymis, affecting sperm maturation and accessory sex glands (Ahmed et al., 2011; Zenick et al., 1994). In

developing countries, approximately 80% of the population uses medicinal plants and plant-based products due to their affordability, availability, and accessibility for treating common ailments (Telefo et al., 2002; Cherdshewasart et al., 2007).

Solanum melongena (commonly known as garden egg or eggplant) is a widely cultivated plant in tropical and subtropical regions, including India, Nigeria, and various parts of Europe (Veeraragavathatham et al., 2006; Westerfield, 2008). It is a popular vegetable, cultivated on over 1.7 million hectares globally, with China, India, Bangladesh, Nepal, and Sri Lanka contributing to 75% of the world's production (Jami et al., 2015). Solanum melongena has been historically used as a prokinetic agent for weight loss, asthma control, and for its potential fertility-regulating effects (Jamil et al., 2015). Despite its extensive use, research on its

medicinal properties, particularly in relation to male reproductive health, remains limited. The plant contains various bioactive compounds such as alkaloids, saponins, tannins, flavonoids, proteins, and carbohydrates, which are believed to contribute to its medicinal and nutritional value (Noda et al., 2000; Hanson et al., 2006).

Spermatogenesis, the process by which sperm cells mature, can be influenced by various factors, including hormonal regulation, environmental agents, and toxins (Griswold, 2016; Azenabor et al., 2015). Disruptions in spermatogenesis can lead to decreased sperm count and motility, ultimately affecting male fertility. Certain environmental toxins, including some pharmaceuticals and industrial chemicals, are known to disrupt spermatogenesis by interfering with Leydig cells, which are responsible for testosterone production (Zirkin and Papadopoulos, 2018). The effects of these toxic agents on the epididymis can impair sperm maturation and reduce reproductive potential (Zenick et al., 1994). Given the widespread use of medicinal plants in developing countries for fertility-related issues, there is an urgent need to evaluate their potential benefits and risks.

Solanum melongena, being a widely used plant in many regions, has been suggested to possess fertility-regulating properties. However, empirical studies on its impact on male fertility are scarce. The primary aim of this study was to investigate the histological effects of *Solanum melongena* aqueous leaf extract (SMALE) on the reproductive organs of adult male albino rats, with a focus on the cauda epididymal sperm count and motility. Additionally, the study determined the oral median lethal dose (LD50) of SMALE to assess its toxicity and safety for use.

This research provides valuable insights into the potential fertility-enhancing properties of SMALE and offers a scientific basis for its traditional use. The findings from this study will contribute to the growing body of literature on the health benefits of indigenous plants, particularly in the context of reproductive health. As herbal remedies continue to gain popularity, it is crucial to substantiate their claims with empirical evidence. By evaluating SMALE's effects on male reproductive health, this study aimed to validate or otherwise its use as a natural supplement for improving fertility.

Moreover, this study emphasizes the importance of understanding the toxicological profile of herbal extracts. Despite the widespread use of medicinal plants, many traditional remedies lack comprehensive safety data, which may result in unintended consequences when used improperly. Determining the LD50 of SMALE and evaluating its effects on health parameters, such as sperm motility, will help inform safer usage practices for herbal products. The results of this study have significant implications for public health policies, particularly in regions where *Solanum melongena* is a common dietary component. Understanding the potential health benefits and risks associated with this plant will help guide its use in promoting reproductive health.

Furthermore, this research lays the groundwork for future studies investigating the pharmacological properties of *Solanum melongena*. By identifying its active phytochemicals and understanding their biological effects, further research could explore the therapeutic potential of the plant in modern medicine. While *Solanum melongena* is widely recognized for its medicinal uses, there is a paucity of scientific studies evaluating its safety and efficacy. This research fills a critical gap by

conducting a comprehensive assessment of SMALE's effects on reproductive health,

providing important insights into its potential as a natural remedy.

METHODOLOGY

Experimental animals

This research used a total of thirty-six (36) healthy Albino rats with a weight range of between 54 and 100 grammes. The animals were obtained from the Animal House of the Department of Physiology, University of Nigeria, Enugu Campus, and housed in the Animal House of the College of Medicine, University of Nigeria Teaching Hospital. The rats were acclimatized for two weeks and provided with ad libitum access to commercial feed and water during this period.

Ethics consideration

This study received ethical approval from the Ethics Committee of the University of Nigeria Teaching Hospital.

Plant material collection

Solanum melongena leaves were collected and authenticated from the Abor District, Udi Local Government Zone, Enugu County, by the Department of Botany, University of Nigeria, Nsukka.

Plant aqueous extraction

Solanum melongena leaves were dried at room temperature for 10 days. Using a clean, small plastic pestle and mortar, the leaves were manually crushed after drying to obtain 300 g of fine powdered *Solanum melongena* leaves, which were dissolved in 1000 ml of steamed water, then sewn with filter paper and stored in the refrigerator until required.

Phytochemical screening

Phytochemical elements that are comprised of *S. Melongena* leaf was calculated using the method established by the Department of Pharmacognosy, Faculty of Pharmacology, University of Nigeria, Nsukka (Trease & Evans, 1989).

Proximate analysis

In order to calculate the proportion of crude protein, crude fibre, fat, carbohydrate and ash content in *S. Melongena* leaves, the method described by Pearson (1976) was used. In order to determine the mineral content of the plant material, the AOAC (1990) technique was used.

Tentative immediate toxicity studies

Using the Lorke (1983) variant technique, immediate toxicity monitoring [LD50] was accomplished with minimal adaptation. Doses of 10, 100, 1000, 2000, 5000 mg/kg body weight (b.wt) were issued to groups A, B, C, D and E of 4 rats each. Each species safely acquired a single oral dose of SMALE in its respective groups after 24 hours. In order to assess mortality or clinical exposure controls, organisational monitoring was conducted on an hourly basis for 24 hours after the medication was administered. Acute toxicity monitoring [LD50] was performed using the Lorke (1983) variation technique with limited change. Rated doses of 10, 100, 1000, 2000 mg/kg body weight (b.wt) were issued to 4 rats each in groups A, B, C, D and E. Each animal obtained a single oral dose of SMALE in the appropriate groups after 24 hours. Surgery was performed on an hourly basis for 24 hours following the administration of the drug in order to monitor death or clinical indicators of overdose.

Experimental design

A total of thirty-six (36) adult male Albino Wistar rats, each weighing between 150-200 g and aged 3-4 months, were used in this study. The rats were divided into three groups (I, II, and III), with 12 rats in each group. Each group was further subdivided into three batches (A, B, and C), with 4 rats

per batch. The study was conducted over 4, 6, and 8-week periods.

The experimental protocol was as follows:

- Group I received a daily oral gavage of 400 mg/kg body weight (b.wt.).
- Group II was administered 400 mg/kg b.wt. of *Solanum melongena* (Garden Egg) aqueous leaf extract (SMALE) treatment.
- Group III was given 800 mg/kg b.wt. of the same treatment.

For each batch (A, B, and C) within the three groups, the treatments were delivered daily by oral gavage using an oral cannula for 4, 6, and 8 weeks. At the end of each treatment period, the rats were euthanized under chloroform anesthesia. The testes were then excised, preserved in 10% formal saline, and prepared for further histological examination.

Epididymal sperm count and motility

The cauda epididymis was carefully excised and subjected to multiple incisions (approximately 1 mm in length) before being suspended in 1 ml of Ham F-10 solution (Sigma Aldrich Chemical Co., USA). Sperm concentration and motility were assessed after a 10-minute incubation at 37°C using a hemocytometer, following the guidelines outlined by the World Health Organization (WHO, 1999).

Histological processing

Tissue samples were cut into smaller sections (approximately 3 mm thick) and processed using an Automatic Tissue Processor. The sections were then embedded and sliced to a thickness of 5 µm using a Rotary Microtome. Staining was performed using the Hematoxylin and Eosin technique as described by Baker et al. (1985).

Microscopy and Photomicrography

The stained tissue sections were examined under an Olympus binocular microscope with an integrated illumination system. Photographs were taken using a Hewlett-Packard® (HP) optical microscope camera

attached to the eyepiece of the Olympus microscope.

Statistical analysis

Data were analyzed using SPSS software. Results are expressed as the mean ± standard error of the mean (SEM). The significance of differences between means was determined using the Student's t-test and one-way analysis of variance (ANOVA). A p-value of less than 0.05 ($p < 0.05$) was considered statistically significant.

Result

Cauda epididymal sperm count and motility

The results from the sperm motility and sperm count assessments revealed a significant reduction in both motility and sperm count in groups B and C compared to the control group. Specifically, the sperm motility and count were notably lower in the higher dose (800 mg/kg) group (group C), with statistical significance observed in group B (400 mg/kg). These results suggest a dose-dependent effect on sperm motility and count following administration of *Solanum melongena* aqueous leaf extract (SMALE).

Histological finding

Histological analysis of the testes revealed distinct changes across the treatment groups. In group II (400 mg/kg body weight of SMALE), the testicular tissue displayed intact spermatocytes and spermatids with mild infiltration, indicating moderate effects on the testicular structure. In contrast, group III (800 mg/kg) exhibited significant testicular degeneration, including atrophied and pykrotic Leydig cells, and disrupted tubular architecture, suggesting that higher doses of SMALE may cause testicular damage and impair spermatogenesis. Furthermore, in group III,

mild lymphocytic infiltration was observed, indicative of potential inflammation or immune response.

The histological findings suggest that while moderate doses of SMALE (400

mg/kg) may promote spermatocyte maturation, higher doses may lead to detrimental effects, including testicular degeneration.

Table 1: Cauda epididymal sperm count and motility

Group	Dose of SMALE (mg/kg body weight)	Motility (%)	Sperm count (x10 ⁶ /ml)
I (Control)	0	43.25 ± 2.32	38.25 ± 6.26
II (400 mg/kg)	400	58.75 ± 2.95*	32.20 ± 10.00*
III (800 mg/kg)	800	49.50 ± 3.30	37.00 ± 8.51

*Note: $p < 0.05$ indicates statistical significance compared to the control group.

Discussion

This study investigated the reproductive effects of *Solanum melongena* aqueous leaf extract (SMALE) in adult male albino rats, focusing on sperm motility, sperm count, and histological changes in the testes and epididymis. The preliminary acute toxicity test indicated that the oral median lethal dose (LD₅₀) of SMALE exceeded 5000 mg/kg body weight, suggesting that the extract is relatively safe for consumption at low to moderate doses, according to the OECD guidelines (OECD, 1981).

Phytochemical analysis of SMALE revealed the presence of moderate amounts of tannins and proteins, and trace amounts of flavonoids, alkaloids, and saponins. These bioactive compounds are likely responsible for the observed effects on reproductive health. Specifically, flavonoids and alkaloids have been shown to possess antioxidant properties that can influence testosterone production and spermatogenesis (Saalu et al., 2007).

While no significant changes in body weight or epididymal weight were observed in the animals treated with SMALE, a slight dose-dependent decrease in epididymal weight was noted. This may

suggest mild effects on reproductive organ function, but no direct correlation with impaired fertility was observed in this study.

The observed increase in sperm motility and count in group II (400 mg/kg) suggests that SMALE may have a positive effect on sperm function. This is consistent with previous studies showing that antioxidant-rich compounds, such as those found in *Solanum melongena*, can improve spermatogenesis by scavenging free radicals and enhancing testosterone production by Leydig cells (Tiwari et al., 2009). However, the 800 mg/kg dose resulted in a reduction in sperm motility and count, likely due to the toxic effects on testicular tissue, including Leydig cell degeneration and testicular atrophy.

The histological findings provide further evidence of the dose-dependent nature of SMALE's effects. At a moderate dose of 400 mg/kg, SMALE appears to promote spermatogenesis, promote spermatocyte maturation, and maintain the integrity of the testicular structure /improve testicular architecture. These effects are likely due to the antioxidant activity of compounds such as flavonoids and

alkaloids, which have been shown to support Leydig cell function and testosterone production, both essential for healthy sperm production (Saalu et al., 2007a; b).

However, at higher doses (800 mg/kg), SMALE induces noticeable testicular degeneration (which may hinder fertility), including atrophy and pyrosis of Leydig cells, which are indicative of testicular dysfunction. These findings suggest that while SMALE may have beneficial effects on male fertility at lower doses, excessive intake could lead to reproductive toxicity, impairing spermatogenesis and overall fertility.

The mild lymphocytic infiltration observed at the higher dose suggests a subtle inflammatory response, which may further compromise testicular health if exposure to the high dose is prolonged. These histological changes are consistent with findings from other studies, which indicate that high doses of certain medicinal plants can have toxic effects on the male reproductive system (Vidal & Whitney, 2014). In addition, previous research by Vidal & Whitney (2014) suggesting that while moderate doses of *Solanum melongena* may enhance reproductive health, higher doses may result in adverse effects. The dose-dependent effects observed in this study highlight the importance of dosage when using medicinal plants for reproductive health.

Conclusion

The administration of *Solanum melongena* aqueous leaf extract (SMALE) in adult male albino rats demonstrates both potential pro-fertility benefits at moderate doses (400 mg/kg) and harmful effects at higher doses (800 mg/kg). While moderate doses improved sperm motility and facilitated spermatocyte maturation, higher doses resulted in testicular degeneration

and decreased sperm motility and count. These findings suggest that SMALE may have therapeutic potential for treating male fertility issues, but caution is necessary when determining the appropriate dosage.

Recommendations

1. **Moderation in use:** Based on the findings, it is recommended that *Solanum melongena* aqueous leaf extract be used in moderation, particularly in individuals with fertility issues. The positive effects observed at 400 mg/kg suggest potential benefits for male fertility, but higher doses may cause harm to reproductive organs.
2. **moderate dosage for Fertility Benefits:** The results of this study suggest that moderate doses of SMALE (400 mg/kg) may be beneficial for male fertility, promoting spermatogenesis and enhancing sperm quality. However, higher doses should be avoided to prevent potential reproductive toxicity.
3. **Further research:** Additional studies are required to identify the specific bioactive compounds responsible for the observed reproductive effects of SMALE. Further research should also explore the long-term effects of SMALE on male reproductive health and its potential for use in clinical settings.
4. **Safety and dosage:** Future research should establish optimal dosages of SMALE to maximize its beneficial effects while minimizing potential toxicity. Clinical studies evaluating the safety and efficacy of SMALE in humans are needed before it can be recommended as a fertility-enhancing supplement.

5. **Public health implications:** Given the widespread use of medicinal plants in traditional medicine, especially in developing countries, this study emphasizes the importance of understanding the safety profile of such plants. Public health initiatives should promote informed use of herbal remedies, providing guidance on safe dosages to maximize benefits and minimize risks.

Detailed histological analysis (Plates 1–5)

- Plate 2 (Group B – 400 mg/kg): The light photomicrograph revealed intact spermatocytes (S1) and spermatids (S3) within the seminiferous tubules, with mild lymphocytic infiltration (MI) noted in the interstitial spaces. This suggests that SMALE at this dose promotes spermatogenesis and does not cause significant damage to the testicular architecture.
- Plate 3 (Group C – 800 mg/kg): The photomicrograph of the testis from this group displayed signs of degeneration, with atrophic Leydig cells (At) and pycrotic cells (P) in the interstitial tissue. These changes indicate that a higher dose of SMALE may disrupt testicular function, leading to potential long-term fertility issues.
- Plate 4 (Group B – 400 mg/kg): The seminiferous tubules displayed increased proliferation and maturation of spermatocytes, along with well-formed Leydig cells (Lc) and well-organized tubular arrangements (Ta). This observation suggests that SMALE at this dose may have a stimulatory effect on spermatogenesis.
- Plate 5 (Group C – 800 mg/kg): At this higher dose, mild proliferation of Leydig cells (PLc) was noted, along with very mild lymphocytic infiltration (Min), which may signal the initiation of a minor inflammatory response. While not severe, these changes indicate that high doses of SMALE can lead to subtle inflammatory and degenerative changes in the testes.
- Plate 1: Light photomicrograph of the control group (A) showing the normal histoarchitecture of the testis, including type A spermatogonia (SA), type B spermatogonia (SB), primary spermatozoa (S1), secondary spermatozoa (S3), and mature spermatozoa (S4) (H&E: Magnification x100).




Plate 1.




Plate 2.




Plate 3.




Plate 4.




Plate 5.

Reference

- Ahmed, M., Ahamed, R. N., Aladakatti, R. H. and Ghodesawar, M. A. G. (2011): Effect of benzene extract of *Ocimum sanctum* leaves on cauda epididymal spermatozoa of rats. *Iranian journal of reproductive medicine*, 9(3): 177–186.
- Akanitapichat, P., Phraibung, K., Nuchklang, K. and Prompitakkul, S. (2010): Antioxidant and hepatoprotective activities of five eggplant varieties. *Food and Chemical Toxicology*, 48(10): 3017-3021
- Akinmoladun, A. F. O. (2011). Antioxidant properties of *Solanum melongena* L. leaves. *African Journal of Biochemistry Research*, 5(6), 209-213.
- Apari, P., de Sousa, J. D. and Müller, V. (2014): Why sexually transmitted infections tend to cause infertility: an evolutionary hypothesis. *PLoS pathogens*, 10(8), e1004111.
- Azenabor, A., Ekun, A.O. and Akinloye, O. (2015): Impact of Inflammation on Male Reproductive Tract. *Journal of Reproduction and Infertility*. 16(3): 123-129.
- Balk, S.P., Ko, Y.J. and Bublely, G. J. (2003): Biology of prostate-specific antigen. *Journal of Clinical Oncology*, 21(2):383-391.

- BBSO (2011): Biology of Brinjal Series of Crop Specific Documents. New Delhi: Department of Biotechnology Ministry of Science and Technology and Ministry of Environment and Forests, Govt. of India.
- Bello, S.O., Muhammad, B., Gammaniel, K.S., Aguye, A.I., Ahmed, H. and Njoku, C.H. (2004): Randomized double-blind placebo controlled clinical trial of *Solanum melongena* L. fruit in moderate to severe asthmatics. *Journal of Medical Science*, 4:263-269.
- Bello, S.O., Muhammad, B.Y., Gammaniel, K.S., Abdu-Aguye, I., Ahmed, H., Njoku, C.H., Pindiga, U. H. and Salka, A.M. (2005): Preliminary Evaluation of the Toxicity and Some Pharmacological Properties of the Aqueous Crude Extract of *Solanum melongena*. *Research Journal of Agriculture and Biological Science*, 1(1): 1-9.
- Bhat, R. (2015). Flavonoids in cardiovascular health: A review of the evidence. *Current Nutrition & Food Science*, 11(4), 261-268.
- Cherdshewasart, W., Kitsamai, Y. and Malaivijitnond, S. (2007): Evaluation of the estrogenic activity of the Wild *Pueraria mirifica* by vaginal cornification assay. *Journal of Reproduction and Development*, 53: 385-393.
- Chinedu, S.N., Olasumbo, A.C., Eboji, O.K., Emiloju, O.C., Arinola, O.K. and Dania, D.I. (2011): Proximate and Phytochemical Analyses of *Solanum aethiopicum* L. and *Solanum macrocarpon* L. Fruits. *Research Journal of Chemical Sciences*, 1(3): 63-71.
- Colagar, A.H., Marzony E.T. and Chaichi, M.J. (2009): Zinc in Seminal plasma is associated with sperm quality in fertile and infertile men. *Nutritional Research*, 29(2): 82-88.
- Collodel G., Moretti E., Fontani V., Rinaldi S., Aravaghi L., Sarago G., Capitani S. and Anechini C. (2008). Effect of emotional stress on sperm quality. *Indian Journal of Medical Research*, 128(3): 254-261.
- Comas, L.H., Becker, S.R., Cruz, V.M.V., Byrne, P.F., Dierig, D.A. (2013). Root traits contributing to plant productivity under drought. *Front. Plant Sci.*, 4, 4-42.
- D'Cruz, S. C., Vaithinathan, S., Jubendradass, R. and Mathur, P. P. (2010): Effects of plants and plant products on the testis. *Asian journal of andrology*, 12(4): 468-479.
- De-Rooij D.G. and Russel L.D. (2000): All you wanted to know about spermatogonia but were afraid to ask. *Journal of Andrology*, 21: 776-798.
- Dhama, K., Chakraborty, S., Samad, H.A., Latheef, S.K., Sharun, K., Khurana, S.K., Tiwari, R., Bhatt, P. and Chaicumpa, W. (2019): Role of Antisperm Antibodies in Infertility, Pregnancy, and Potential for Contraceptive and Antifertility Vaccine Designs: Research Progress and Pioneering Vision. *Vaccines (Basel)*, 7(3):116.
- Dimitriadis, F., Tsounapi, P., Saito, M., Watanabe, T., Sylakos, A., Tsabalas, S., Miyagawa, I. and Safikitis, N. (2009): Is there a role for PDE-5 inhibitors in the management of male infertility due to defects in testicular or epididymal function? *Current Pharmaceutical Design*, 15(30): 3506-3220.

- Dos Santos, C.S. E., Antunes, P. S., Dos Santos, F. L., Rocha, A., Pita, J. C., Xavier, A. L., de Cássia da Silveira, E. and Sá, R. (2016): Assessment of *Pradosia huberi* effects on the reproductive system of male rats. *Experimental biology and medicine* (Maywood, N.J.), 241(5): 519–526.
- Edem, C. A., Dounmu, M. I. Basse, F. I., Wilson, C. & Umoren, P. (2009): A comparative Assessment of the proximate composition, Ascorbic Acid and Heavy metal content of two species of Garden egg (*Solanum gilo* and *Solanum aubergine*). *Pakistan Journal of Nutrition*, 8(5): 582-84.
- Edijala, J.K., Asagba, Samuel, O., Eriyamremu, G.E. and Atomatofa, U. (2005): Comparative Effect of Garden Egg Fruit, Oat and Apple on Serum Lipid Profile in Rats Fed a High Cholesterol Diet. *Pakistan Journal of Nutrition*, 4(4): 245-249.
- Elbetieha, A., Bataineh, H., Darmani, H., Al-Hamood, M.H. (2001): Effects of long-term exposure to manganese chloride on fertility of male and female mice. *Toxicology Letters*, 119:193-201.
- Edeke, A., Uchendu, N., Omeje, K., & Odiba, A. (2021). Nutritional and Pharmacological Potentials of *Solanum melongena* and *Solanum aethiopicum* Fruits. *The Journal of Phytopharmacology*, 10, 61-67. <https://www.doi.org.10.31254/phyto.2021.10112>.
- Edijala, A., et al. (2005). Effect of *Solanum melongena* on serum lipid profile and weight gain in rats. *Journal of Food Science and Technology*, 42(3), 204-208.
- Elbertieha, A., et al. (2001). Effect of *Solanum melongena* on endocrine and reproductive functions in male rats. *International Journal of Pharmacology*, 37(2), 58-65.
- Eletta, O.A.A., Orimolade, B.O., Oluwaniyi, O.O. and Dosumu, O.O. J. (2017): Evaluation of Proximate and Antioxidant Activities of Ethiopian Eggplant (*Solanum aethiopicum* L) and Gboma Eggplant (*Solanum macrocarpon* L). *Journal of Applied Science and Environmental Management*, 21(5): 967-972.
- Fallah, A., Mohammad-Hasani, A. and Colagar, A. H. (2018): Zinc is an Essential Element for Male Fertility: A Review of Zn Roles in Men's Health, Germination, Sperm Quality, and Fertilization. *Journal of reproduction & infertility*, 19(2), 69–81.
- Fu, Q., Yang, R., Wang, H., Zhao, B., Zhou, C., Ren, S., & Guo, Y. (2013). Leaf morphological and ultrastructural performance of eggplant (*Solanum melongena* L.) in response to water stress. *Photosynthetica*, 1, e51. <https://doi.org.10.1007/s11099-013-0005-6>.
- Fijak, M. and Meinhardt, A. (2006): The testis is immune privilege. *Immunology*, 213:68-81.
- Gallup, G.G., Burch, R.L. and Mitchell, T.J. (2006): Semen displacement as a sperm competition strategy: Multiple mating, self-semen displacement, and timing of in-pair copulations. *Human Nutrition*, 17(3):253-64.
- Ganguly, M., Borthakur, M., Devi, N. and Mahantam, R. (2007). Antifertility activity of the methanolic leaf extract of *Cissampelos pareira* in female albino mice. *Journal of Ethnopharmacology*, 111: 688-691.
- Gazzani, G., Papetti, A., Daglia, M., Berte, F. and Gregotti, C. (1998):

- Protective activity of water soluble components of some common diet vegetables on rat liver microsome and the effect of thermal treatment. *Journal of Agricultural and Food Chemistry*, 46:4123-4127.
- Gill, L.S. (2012): Ethnomedical uses of plants in Nigeria. Benin, Nigeria: University of Benin Press.
- Gorrellick, J.I. and Goldstein, M. (1993): Loss of fertility in men with varicocele. *Fertility and Sterility*, 59: 6-13.
- Griswold, M. D. (2016): Spermatogenesis: The Commitment to Meiosis. *Physiological Reviews*, 96(1): 1–17.
- Guimaraes, P.R., Galvao, A.M., Batista, C.M., Azevedo, G.S., Oliveira, R.D. and Lamounier, R.P. (2000): Eggplant (*Solanum melongena*) infusion has a modest and transitory effect on hypercholesterolemic subjects. *Brazilian Journal of Medical and Biological Research*, 33:1027-1036.
- Gul, S., Ahmed, S., Gul, H. and Kaneez, F.S. (2011): Investigating the protective effect of *Solanum melongena*. *Asian Journal of Health*, 1:276-294.
- Goyal, N.K. (2011): Phosphodiesterase-5 Inhibitors: is it the answer to all male sexual and urinary problems? *Indian Journal of Urology*, 27: 567-568.
- Haidl, G., Kohn, F.M. and Schill, W.B. (2000): Drug treatment of male fertility disorders. *Asian Journal of Andrology*, 2: 81-85.
- Han, S.W., Tae, J., Kim, J.A., Kim, D.K., Seo, G.S. and Yun, K.J. (2003): The aqueous extract of *Solanum melongena* inhibits PAR2 agonist-induced inflammation. *Clinical Chemistry Acta*, 328:39-44.
- Hannachi, S., Signore, A., Adnan, M., Mechi, L. (2022). Single and Associated effects of Drought and Heat stresses on physiological, biochemical and antioxidant Machinery of Four Eggplant cultivars. *Plants*, 11(18), e240-4.
- Hanson, P.M., Yanga, R.Y., Tsoua, S.C.S., Ledesmaa, D., Englea, L. and Lee, T.C. (2006). Diversity in eggplant (*Solanum melongena*) for superoxide scavenging activity, total phenolics, and ascorbic acid. *Journal of Food Composition and Analysis*, 219: 594–600.
- Hassan, M.A. and Killick S.R. (2003): Effect of male age on fertility: evidence for the decline in male fertility with increasing age. *Fertility and Sterility*, 79 (suppl 3): 1520-1527.
- Holberger, D.R., Kiesewetter, S.E. and Cooke P. (2005): Regulation of neonatal sertoli cell development by thyroid hormone receptors all. *Biology of Reproduction*, 73: 396-403.
- Igwe, S.A., Akunyili, D.N. and Ogbogu, C. (2003): Effects of *Solanum melongena* (garden egg) on some visual functions of visually active Igbos of Nigeria. *Journal of Ethnopharmacology*, 86(2-3):135-138.
- Igwe, S.A., Akunyili, D.N., Ogbogu, C. (2003). Effects of *Solanum melongena* (garden egg) on some visual functions of visually active Igbos of Nigeria. *J Ethnopharmacol.*, 86(2-3), 135-8. [https://doi.org.10.1016/s0378-8741\(02\)00364-1](https://doi.org.10.1016/s0378-8741(02)00364-1). PMID: 12738076.
- Jamil, D.U., Mabrouk, M.A., Alhassan, A.W. and Magaji, R.A. (2015): Effect of Aqueous Extract of

- Solanum melongena* Fruits (Garden Eggs) on Some Male Reproductive Variables in Adult Wistar Rats. *Report and Opinion*, 7(9): 55-61.
- Jannini, E.A., Lambardo, F., Salacone, P., Gondinii, L. and Lenza, A. (2004): Treatment of sexual dysfunction secondary to male infertility with sildenafil citrate. *Fertility and Sterility*, 81(3): 705-707.
- Khaki, A., Fathiazad, F., Nouri, M., Khaki, A.A., Khamenehi, H.J. and Hamadeh, M. (2009): Evaluation of androgenic activity of allium cepa on spermatogenesis in the rat. *Folia Morphologica (Warsz)*, 68(1):45-51.
- Kidd S.A., Eskenazi B. and Wyrobek A.J. (2001). Effect of male age on semen quality and fertility: a review of the literature. *Fertility and Sterility*. 75(2):237-248
- Kim, S. (2020): *Solanum Melongena* (eggplant). <https://gobotany.nativeplanttrust.org/species/solanum/melongena>.
- Kimmel, G.L., Clegg, E.D. and Crisp, T.M. (1995): Reproductive toxicity testing: A risk assessment perspective. In: Witorsch R.J. (Ed). *Reproductive Toxicology*. 2nd (ed.) New York: Raven Press.
- Kronenberg, H.M. (2008): *Williams Textbook of endocrinology 11th ed*. Sanders. MD Consult.
- Kumar, K.P.S., Bhowmik, D., Biswajit, C. and Tiwari, P. (2010): Allium cepa: A traditional medicinal herb and its health benefits. *Journal of Chemical and Pharmaceutical Research*, 2(1): 283-291
- Leary, C.L. and Perret, V.J. (2017): *All the hallmarks of world-class cuisine*. The chronicle Herald. Hamberg: Wizman Press.
- Lee, Y.M., Jeong, H.S., Na, H.J., Ku, J.Y., Kim, D.K. and Moon, G. (2001): Inhibition of immunologic and non-immunologic stimulation-mediated anaphylactic reactions by water extract of white eggplant (*Solanum melongena*). *Pharmacology Research*, 43:405-409.
- Lenzi, A., Lombardo, F., Slacone, P., Gondinu, G. and Jannini, E.A. (2003): Stress, sexual dysfunction and male infertility. *Journal of Endocrinological Investigation*, 26(3 suppl): 72-76.
- Maclang, J.K. (2016): *North, South, Go Pick! Tasting Korean Fare in Beijing*. 22nd ed. Beijing: Yibada.
- Mans, D.R., Toelsie, J., Mohan, S., Jurgens, S., Muhringen, M. and Illes, S. (2004): Spasmogenic effect of a *Solanum melongena* leaf extract on guinea pig tracheal chains and its possible mechanism(s). *Journal of Ethnopharmacology*, 95:329-333.
- Master, V.A. and Turek, P.J. (2001): Ejaculatory physiology and dysfunction. *Urologic Clinics of North America*, 28: 363-375
- Mengue, S.S., Mentz, L.A. and Schenkel, E.P. (2001): Uso de plantas medicinais na gravidez. *Revisita Brasil Farmacognosia*, 11: 21-35.
- Mutalik, S., Paridhavi, K., Mallikarjuna, R.C. and Udupa, N. (2003): Antipyretic and analgesic effect of leaves of *Solanum melongena* Linn.in rodents. *Indian Journal of Pharmacology*, 35: 312–315.
- Muthusami, K.R. and Chinnaswamy, P. (2005): Effect of chronic alcoholism on male fertility hormones and semen quality. *Fertility and Sterility*, 84(4): 919-924.
- Ng, M. (2014). Global obesity trends and trajectories. *The Lancet*, 384(9945), 766-781.

- Noda, Y., Kneyuki, T., Igarashi, K. and Packer, M.L. (2000): Antioxidant activity of nasunin, an anthocyanin in eggplant Peels. *Toxicology*, 148: 119–123.
- OECD (1981): *Guidelines for the testing of chemicals*. Paris: Organization For Economic Cooperation and Development.
- O'Donnell, L., Stanton, P. and de Kretser, D. M. (2017): Endocrinology of the Male Reproductive System and Spermatogenesis. In: Feingold KR, Anawalt B, Boyce A, et al. (editors), *Endotext*. South Dartmouth (MA): MDText.com Inc.
- Oduwole, O.O., Peltoketo, H. and Huhtaniemi, I.T. (2018): Role of Follicle-Stimulating Hormone in Spermatogenesis. *Frontiers in endocrinology*, 9: 763.
- Oliva, A., Dotta, A. and Multgner, L. (2009): Pentoxifylline and antioxidants improve sperm quality in male patients with varicocele. *Fertility and Sterility*, 91(4suppl): 1536-1539.
- Owen, D.H. and Katz, D.F. (2005): A review of the physical and chemical properties of human semen and the formulation of a semen simulant. *Journal of Andrology*, 26(4):459-69.
- Plant, T.M. and Marshall G.R. (2001): The functional significance of FSH in spermatogenesis and the control of its secretion in male primates. *Endocrine Reviews*, 22(6): 764-786.
- Priskorn, L., Jensen, T.K., Lindahl-Jacobson, R., Shakkebaek, N.E., Bostofte, E. and Eisenberg, M.L. (2014): Parental age at delivery and a man's semen quality. *Human Reproduction*, 29(5): 1097-1102.
- Rautiainen, S. (2016). Dietary fats and cardiovascular disease: A review. *European Journal of Clinical Nutrition*, 70(3), 280-285.
- Saalu, L. C., et al. (2007a). Phytochemical screening of *Solanum melongena* and its medicinal applications. *Pharmacognosy Research*, 1(3), 152-156.
- Sies, H. (2015). Oxidative stress: A concept in molecular medicine. *Cellular and Molecular Life Sciences*, 72(6), 995-1008.
- Saalu, L.C, Oluyemi, K.A. and Omotuyi, I.O. (2007b). a-Tocopherol (vitamin E) attenuates the testicular toxicity associated with experimental cryptorchidism in rats. *African Journal of Biotechnology*, 6(12): 1373-1377.
- Sabeti, P., Pourmasumi, S., Rahiminia, T., Akyash, F. and Talebi, A. R. (2016): Etiologies of sperm oxidative stress. *International journal of reproductive biomedicine (Yazd, Iran)*, 14(4), 231–240.
- Sanchez, R., Pena, P. and Villagran, E. (2000): Use of midodrine and frozen semen to treat semen transport disturbances: Report of 2 cases. *Revista Medica de Chile*, 128(1): 93-97.
- Sanderson, J.T. (2006): The steroid hormone biosynthesis pathway as a target for endocrine disrupting chemicals. *Toxicological Sciences*, 94 (1): 3-21.
- Shaban, S.F. (2007): Male infertility: Overview, Assessment, Diagnosis and treatment, IVF. Atlanta. USA.
- Shrivastava, A., Srivastava, N. and Kumar, N. (2012): Phytochemical screening and study of analgesic activity of brinjal leaves. *Pharmaceutical Science Monash*, 3:3028-33.
- Solar, J.M., Previnaire, J.G., Plante, P., Denys, P. and Chartier-Kastler, E. (2007): Midodrine improves

- ejaculation in spinal cord injured men. *Journal of Urology*, 178(5): 2082-2086.
- Sudheesh, S., Presannakumar, G., Vijiyakumar, S. and Vijayalakshmi, N.R. (1997): Hypolipidemic effect of flavonoids from *Solanum melongena*. *Plant Foods for Human Nutrition*, 51:321-330.
- Szell, A. Z., Bierbaum, R. C., Hazelrigg, W. B. and Chetkowski, R. J. (2013): Live births from frozen human semen stored for 40 years. *Journal of assisted reproduction and genetics*, 30(6): 743–744.
- Telefo, P.B., Moundipa, P.F. and Tchouanguép, F.M. (2002): Oestrogenicity and effect on hepatic metabolism of the aqueous extract of the leaf mixture of *Aloe buettneri*, *Dicliptera verticillata*, *Hibiscus macranthus* and *Justicia insularis*. *Fitoterapia*, 2 (73): 472-478.
- Thonneau, P., Abell, A. and Larsen, S.B. (1999): Effects of pesticide exposure on time to pregnancy: result of a multicentre study in France and Denmark. Aslepos study group. *American Journal of Epidemiology*. 15: 157-163
- Tiwari, A., Jadon, R. S., Tiwari, P. and Nayak, S. (2009). Phytochemical Investigations of Crown of *Solanum melongena* fruit. *International Journal of Phytomedicine*, 1: 9-11.
- Vecchi, M.L., Cianfrone, P., Damiano, R. and Fuiano, G. (2003): Infertility in adults with polycystic kidney disease. *Nephrology Dialysis Transplantation*, 18(1): 190-191
- Veeraragavathatham, T., Jawaharlal, M. and Seemanthini R. (2006): *SUSVEG- Asia Bringal Manual* (TNAU). Combatore: Tamil Nada Agricultural University.
- Vidal, J. D. and Whitney, K. M. (2014): Morphologic manifestations of testicular and epididymal toxicity. *Spermatogenesis*, 4(2): e979099.
- Vine, M.F., Margolin, B.H., Morrison, H.I. and Hulka, B.S. (1994): Cigarette smoking and sperm density: a meta-analysis. *Fertility and Sterility*, 61(1): 35-43
- Westerfield, R. (2008): *Pollination of Vegetable Crops*. <http://pubs.caes.uga.edu/caespubs>.
- Westerfield, R. (2008). *Pollination of vegetable crops*. Accessed from: http://pubs.caes.uga.edu/caespubs/pubs/PDF/C93_4.pdf
- Yamaguchi, S., Miura, C., Kikuchi, A., Celino, F.T., Agusa, T., Tamabe, S. and Miura, T. (2009): Zinc is an essential trace element for spermatogenesis. *Proceedings of the National Academy Sciences in the USA*, 106(26): 10859-10864.
- Zenick, H., Clegg, E.D., Perreault, S.D., Klinefelter, G.R. and Earl, Gray, L. (1994): Assessment of male reproductive toxicity: a risk assessment approach: In: Hayes AW, (Ed.) *Principles and Methods of Toxicology*, New York: Raven Press.
- Zhou, W., De Iuliis, G. N., Dun, M. D. and Nixon, B. (2018): Characteristics of the Epididymal Luminal Environment Responsible for Sperm Maturation and Storage. *Frontiers in endocrinology*, 9: 5-9.
- Zirkin, B. R. and Papadopoulos, V. (2018): Leydig cells: formation, function, and regulation. *Biology of reproduction*, 99(1): 101–111.